arXiv:math/0601513v1 [math.OA] 20 Jan 2006

Tracial Rokhlin property for automorphisms on simple $A\mathbb{T}$-algebras

Huaxin Lin
Department of Mathematics
University of Oregon
Eugene, Oregon 97403-1222
Hiroyuki Osaka
Department of Mathematical Sciences
Ritsumeikan University
Kusatsu, Shiga, 525-8577, Japan

Abstract

Let A be a unital simple $A\mathbb{T}$-algebra of real rank zero. Given an isomorphism $\gamma_1 : K_1(A) \to K_1(A)$, we show that there is an automorphism $\alpha : A \to A$ such that $\alpha_{*1} = \gamma_1$ which has the tracial Rokhlin property. Consequently, the crossed product $A \rtimes_\alpha \mathbb{Z}$ is a simple unital AH-algebra with real rank zero. We also show that automorphism with Rokhlin property can be constructed from minimal homeomorphisms on a connected compact metric space.

1 Introduction

Alan Connes introduced the Rokhlin property in ergodic theory to operator algebras ([2]). Several versions of the Rokhlin property for automorphisms on C^*-algebras have been studied (for example [8], [32], [14], [27], [11], [12] and [30], to name a few). Given a unital C^*-algebra A and an automorphism α on A, one may view the pair (A, α) as a non-commutative dynamical system. To study its dynamical structure, it is natural to introduce the notion of Rokhlin property. Let A be a unital simple $A\mathbb{T}$-algebra (a C^*-algebra which is an inductive limit of those C^*-algebras that are finite direct sums of continuous functions on the circle $\mathbb{T}$) and let α be an automorphism on A. In several cases, Kishimoto showed that if α is approximately inner (or homotopic to the identity) and has a Rokhlin type property, then the crossed product $A \rtimes_\alpha \mathbb{Z}$ is again a unital simple $A\mathbb{T}$ of real rank zero ([14], [15], [16]). It is proved that if α induces the identity on $K_0(A)$ (or a "dense" subgroup of $K_0(A)$) and α has so-called tracial cyclic Rokhlin property then indeed the crossed product is an AH-algebra of real rank zero ([25] and [24]). A natural question is when automorphisms have certain Rokhlin property. In [29], it is shown that if A is a unital separable simple C^*-algebra with tracial rank zero and with a unique tracial state and if α is an automorphism such that $A \rtimes_\alpha \mathbb{Z}$ has a unqiue trace, then α has the tracial Rokhlin property. More recently, N.C. Phillips [31] showed that, for any unital simple separable C^*-algebra A with tracial rank zero, there is a dense G-δ set of approximately inner automorphisms such that every automorphism in the the set has the tracial Rokhlin property.

Let A be a unital simple $A\mathbb{T}$-algebra of real rank zero. Suppose that $\gamma_1 : K_1(A) \to K_1(A)$ is an automorphism. In this note, we present an automorphism α on A with the tracial cyclic Rokhlin property such that $\alpha_{*1} = \gamma_1$. The automorphism α that we present in this note also has the property that $\alpha_{*0} = \mathrm{id}_{K_0(A)}$. When α has the tracial cyclic Rokhlin property, then α must fix a large subgroup of $K_0(A)$. In fact, it is shown in [24] that, at least in the case that A has a unique tracial state, if α has tracial cyclic Rokhlin property, then α^2 fixes a subgroup $G \subset K_0(A)$ so that $\rho(G)$ is dense in $Aff(T(A))$, where $T(A)$ is the tracial state space of A, $Aff(T(A))$ is the space of all real affine continuous functions on $T(A)$ and $\rho : K_0(A) \to Aff(T(A))$ is the positive homomorphism induced by the evaluation $\rho([p])(\tau) = \tau(p)$ for projections $p \in A$.

Let X be a connected compact metric space and $\rho_X : K_0(C(X)) \to \mathbb{Z}$ be the dimension map. Denote by $\mathrm{ker}\rho_X$ the kernel of the dimension map. Given a such X and a countable dense subgroup $D \subset \mathbb{Q}$, there is a standard way to construct a unital simple C^*-algebra A_X with tracial rank zero such that $K_0(A_X) = D \oplus \mathrm{ker}\rho_X$ and $K_1(A_X) = K_1(C(X))$ as well as there is a unital embedding $j : C(X) \to A_X$ so that $(j_{*0})|_{\mathrm{ker}\rho_X} = \mathrm{id}|_{\mathrm{ker}\rho_X}$ and $j_{*1} = \mathrm{id}_{K_1(X)}$. This could be viewed as a version of the non-commutative space associated with X. Suppose that $\psi : X \to X$ is a minimal homeomorphism on X. We show that one can construct an automorphism α on A_X associated with ψ such that α has the tracial cyclic Rokhlin property and such that $\alpha_{*0}|_D = \mathrm{id}_D$, $\alpha_{*0}|_{\mathrm{ker}\rho_X} = \psi_{*0}|_{\mathrm{ker}\rho_X}$ and $\alpha_{*1} = \psi_{*1}$. This may be viewed as a non-commutative version of the minimal action associated with ψ. We also show that somewhat general construction can also be made. It appears that automorphisms with the tracial cyclic Rokhlin property occur quite often. While we believe that many other types of construction of automorphisms with the Rokhlin property may be possible and perhaps not necessarily difficult, we think these construction in this note shed some light on how commutative Rokhlin tower lemma appears naturally in the study of non-commutative dynamical systems such as (A, α).

Aknowledgement The first named author was partially supported by a NSF grant. The second author was partially supported by Open Research Center Project for Private Universities:maching fund from MEXT,2004-2008. Much of the ground work of this reseach was done when both authors were visiting East China Normal University in the summer 2004. They would like to acknowledge the support from Shanghai Priority Academic Disciplines and from Department of Mathematics of East China Normal University.

2 Preliminaries

The following conventions will be used in this paper.

(1) Let A be a stably finite C^*-algebra. Denote by $T(A)$ the tracial state space of A. Denote by $Aff(T(A))$ the normed space of all real affine continuous functions on $T(A)$.

(2) Denote by $\rho_A : K_0(A) \to Aff(T(A))$ the positive homomorphism induced by $\rho_A([p])(\tau) = \tau \otimes Tr(p)$ for any projection in $M_k(A)$ $(k = 1, 2, ...,)$, where Tr is the standard trace on M_k.

(3) Let X be a connected compact metric space. Denote by $\rho_X : K_0(C(X)) \to \mathbb{Z}$ the positive homomorphism $\rho_{C(X)}$ defined in (2). It is the dimension function from $K_0(C(X))$ to $\mathbb{Z}$.

(4) Let X be a compact metric space, let F be a subset of X and let $\varepsilon > 0$. Put

$$F_\varepsilon = \{x \in X : \mathrm{dist}(x, F) < \varepsilon\}.$$

(5) Let A be a C^*-algebra and let p and q be projections in A. We say p is equivalent to q if there is a $v \in A$ such that $v^*v = p$ and $vv^* = q$.

(6) Let $A = \lim_{n\to\infty}(A_n, \phi_n)$ be an inductive limit of C^*-algebras. Here ϕ_n is a homomorphism from A_n into A_{n+1}. We will use $\phi_{n,\infty} : A_n \to A$ for the homomorphism induced by the inductive limit system. We also use $\phi_{n,m}$ for $\phi_{m-1} \circ \phi_{m-2} \circ \cdots \circ \phi_n$, if $m > n$. Suppose that each $A_n = \oplus_{i=1}^{r(n)} B_{n,i}$. When $B_{n,i}$ are understood, a partial map $\phi_n^{(i,j)}$ of ϕ_n is the homomorphism from $B_{n,i}$ to $B_{n+1,j}$ given by ϕ_n.

(7) Let A and B be two C^*-algebras and let $\phi, \psi : A \to B$ be two maps. Suppose that $\mathcal{G} \subset A$ is a subset. We write

$$\psi \approx_\varepsilon \phi \ \text{ on } \ \mathcal{G},$$

if

$$\|\phi(a) - \psi(a)\| < \varepsilon \ \text{ for all } \ a \in \mathcal{G}.$$

The following Rokhlin property was introduced in [28].

Definition 2.1. Let A be a unital C^*-algebra and let α be an automorphism on A. We say that α has the tracial Rokhlin property if, for any $\varepsilon > 0$, any finite subset $\mathcal{F} \subset A$, any positive integer $n > 0$, and any $a \in A_+ \setminus \{0\}$, there are mutually orthogonal projections $e_1, e_2, ..., e_n$ such that

(1) $\|e_i x - x e_i\| < \varepsilon$ for all $x \in \mathcal{F}$, $(0 \le i \le n-1)$

(2) $\|\alpha(e_i) - e_{i+1}\| < \varepsilon$, $i = 1, 2, ..., n-1$ and

(3) $1 - \sum_{i=0}^{n-1} e_i$ is equivalent to a projection in $\overline{aAa}$.

A stronger version of the Rokhlin property below was given in [25].

Definition 2.2. Let A be a unital C^*-algebra and let α be an automorphism on A. We say that α has the tracial *cyclic* Rokhlin property if, for any $\varepsilon > 0$, any finite subset $\mathcal{F} \subset A$, any positive integer $n > 0$, and any $a \in A_+ \setminus \{0\}$, there are mutually orthogonal projections $e_0, e_1, e_2, ..., e_{n-1}$ (with $e_n = e_0$) such that

(1) $\|e_i x - x e_i\| < \varepsilon$ for all $x \in \mathcal{F}$, $(0 \le i \le n-1)$

(2) $\|\alpha(e_i) - e_{i+1}\| < \varepsilon$, $i = 0, 1, ..., n-1$ and

(3) $1 - \sum_{i=0}^{n-1} e_i$ is equivalent to a projection in $\overline{aAa}$.

Remark 2.3. Note that, in Definition 2.2, one requires that $\|\alpha(e_{n-1}) - e_0\| < \varepsilon$. This is not required in Definition 2.1. If A is assumed to satisfy the so-called fundamental comparison property, i.e., $\tau(p) > \tau(q)$ for all $\tau \in T(A)$ implies that q is equivalent to a projection $e \le p$ for any pair of projections, then condition (3) above can be replaced by $\tau(1 - \sum_{i=0}^{n-1} e_i) < \varepsilon$ for all $\tau \in T(A)$ and the reference to the element a can be removed. We note that if A has tracial rank zero then A has the fundamental comparison property.

If A has tracial rank zero, then A has real rank zero, stable rank one, and $\rho_A(K_0(A))$ is dense in $Aff(T(A))$. One of the important consequences that a given automorphism satisfies the tracial cyclic Rokhlin property is the following result:

Theorem 2.4. (see 3.4 and 4.5 of [24])

If A is a unital separable amenable simple C^*-algebra with tracial rank zero satisfying the UCT and α is an automorphism on A with the tracial (cyclic) Rokhlin property such that $(\alpha)_{*0} = \mathrm{id}_G$ for some subgroup $G \subset K_0(A)$ for which $\rho_A(G)$ is dense in $\rho_A(K_0(A))$, then $A \rtimes_\alpha \mathbb{Z}$ has tracial rank zero.

Consequently, $A \rtimes_\alpha \mathbb{Z}$ is an AH-algebra.

3 Automorphisms on simple $A\mathbb{T}$-algebras

The purpose of this section is to present Theorem 3.5.

We start with the following construction which is certainly familiar to experts.

Lemma 3.1. *Let A be a unital simple infinite dimensional AF-algebra. Then, there exists a sequence of finite dimensional C^*-subalgebras $F_n = M_{k(n)} \oplus B_n$ and monomorphisms $\phi_n : F_n \to F_{n+1}$ satisfying the following:*

(1) *Let d_n be the identity of $M_{k(n)}$. Then*

$$\lim_{n\to\infty} \sup_{\tau\in T(A)} \{\tau(\phi_{n,\infty}(d_n))\} = 0;$$

(2) *The partial map from $M_{k(n)}$ to $M_{k(n+1)}$ has multiplicity at least 2.*

(3) $A = \lim_{n\to\infty}(F_n, \phi_n)$.

Proof. We first write $A = \lim_{n\to\infty}(A_n, \psi_n)$, where each A_n is a finite dimensional C^*-algebra and ψ_n is a monomorphism.

Write $A_1 = M_{n(1)} \oplus B_1'$. Since A is assumed to be simple, we may also assume that the partial map $\psi_n^{1,1}$ from $M_{n(1)}$ to a simple summand $M_{n(2)}$ of A_2 has multiplicity $m(1)$ which is at least 4. Define $F_1 = M_{n(1)} \oplus M_{n(1)} \oplus \cdots \oplus M_{n(1)} \oplus B_1'$, where $M_{n(1)}$ repeats $[m(1)/2]$ (integer part of $m(1)/2$) times. Denote $B_1 = M_{n(1)} \oplus \cdots \oplus M_{n(1)} \oplus B_1'$, where $M_{n(1)}$ repeats $[m(1)/2] - 1$ times. So $F_1 = M_{n(1)} \oplus B_1$. Note there are monomorphism $f_1 : A_1 \to F_1$ and monomorphism $\psi_1' : F_1 \to A_2$ such that $\psi_1 = \psi_1' \circ f_1$ and the multiplicity of the partial map of ψ_1' from $M_{n(1)}$ to $M_{n(2)}$ is at least 2. Let d_1 be the identity of $M_{n(1)}$ in F_1. Note that

$$\tau(\psi_{2,\infty} \circ \psi_1'(d_1)) \le 1/2$$

for all $\tau \in T(A)$.

We have $A_2 = M_{n(2)} \oplus B_2'$, where B_2' is a finite dimensional C^*-algebra.

We may also assume (by replacing A_3 by some A_n) that the partial map of ψ_2 from $M_{n(2)}$ to a simple summand $M_{n(3)}$ of A_3 has multiplicity $m(2)$ which is at least least 8. Define $F_2 = M_{n(2)} \oplus M_{n(2)} \oplus \cdots M_{n(2)} \oplus B_2'$, where $M_{n(2)}$ repeats $[m(2)/2]$ times. Define $B_2 = M_{n(2)} \oplus \cdots \oplus M_{n(2)} \oplus B_2'$, where $M_{n(2)}$ repeats $[m(2)/2] - 1$ times. So $F_2 = M_{n(2)} \oplus B_2$. There exists a monomorphism $f_2 : A_2 \to F_2$ and there exists a monomorphism $\psi_2' : F_2 \to A_3$ such that

$$\psi_2 = \psi_2' \circ f_2.$$

Moreover, the multiplicity of the partial map of ψ_2' from $M_{n(2)}$ to $M_{n(3)}$ is at least 2.

Define $\phi_1 : F_1 \to F_2$ by $\phi_1 = f_2 \circ \psi_1'$. Let d_2 be the identity of the first summand $M_{n(2)}$ in F_2. Then

$$\tau(\psi_{3,\infty} \circ \psi_2'(d_2)) < 1/4$$

for all $\tau \in T(A)$.

Write $A_3 = M_{n(3)} \oplus B_3'$, where B_3' is a finite dimensional C^*-subalgebra. We may assume that the partial map of ψ_3 from $M_{n(3)}$ to a simple summand $M_{n(4)}$ of A_4 has multiplicity $m(3)$ at least 16. Define $F_3 = M_{n(3)} \oplus M_{n(3)} \oplus \cdots \oplus M_{n(3)} \oplus B_3'$, where $M_{n(3)}$ repeats $[m(3)/2]$ times and $B_3 = M_{n(3)} \oplus \cdots \oplus M_{n(3)} \oplus B_3'$, where $M_{n(3)}$ repeats $[m(3)/2] - 1$ times. So $F_3 = M_{n(3)} \oplus B_3$.

There are monomorphisms $f_3 : A_3 \to F_3$ and $\psi_3' : F_3 \to A_4$ such that $\psi_3 = \psi_3' \circ f_3$. Moreover, the partial map of ψ_3 from $M_{n(3)}$ (the first summand) to $M_{n(4)}$ has multiplicity at least 2. Define $\phi_2 : F_2 \to F_3$ by $\phi_2 = f_3 \circ \psi_2'$. Let d_3 be the identity of of the first summand $M_{n(3)}$ in F_3. Then

$$\tau(\psi_{4,\infty} \circ \psi_3'(d_3)) < 1/8.$$

We continue this construction. We obtain $F_k = M_{n(k)} \oplus B_k$, where B_k is a finite dimensional C^*-subalgebra, and we obtain monomorphisms $f_k : A_k \to F_k$, $\psi'_k : F_k \to A_{k+1}$ and $\phi_k : F_k \to F_{k+1}$ such that

$$\psi_k = \psi'_k \circ f_k \quad \text{and} \quad \phi_k = f_{k+1} \circ \psi'_k \tag{e 3.1}$$

and such that the partial map of ϕ_k from $M_{n(k)}$ to $M_{n(k+1)}$ has multiplicity at least 2. Moreover,

$$\tau(\psi_{k+1,\infty} \circ \psi'_k(d_k)) < 1/2^k \tag{e 3.2}$$

for all $\tau \in T(A)$, where d_k is the identity of $M_{n(k)}$ in F_k.

By (e 3.1), $A = \lim_{n\to\infty}(F_k, \phi_k)$. By (e 3.2),

$$\tau(\phi_{k,\infty}(d_k)) < 1/2^k.$$

The lemma follows.

□

Lemma 3.2. *Let $X = S^1 \vee S^1 \vee \cdots \vee S^1$ be identified with m copies of unit circle with a common point 1. For any integer n and finite subset $\mathcal{F} \subset C(X)$, there exists an integer $N = \Lambda(n, \mathcal{F})$ satisfying the following:*

For any unital C^*-algebra B with real rank zero and any homomorphisms $\phi_1, \phi_2 : C(X) \to B$ with $(\phi_1)_{*1} = (\phi_2)_{*1}$, there is a unitary $u \in M_{mN+1}(B)$ such that

$$u^*(\mathrm{diag}(\phi_1(f), f(\xi_1), f(\xi_2), ..., f(\xi_{mN}))u \approx_{1/2^n}$$

$$\mathrm{diag}(\phi_2(f), f(\xi_1), f(\xi_2), ..., f(\xi_{mN})),$$

where each of $\{\xi_1, \xi_2, ..., \xi_N\}$, $\{\xi_{N+1}, \xi_{N+2}, ..., \xi_{2N}\}$, ..., $\{\xi_{(m-1)N+1}, \xi_{(m-1)N+2}, ..., \xi_{mN}\}$, divides the unit circles evenly.

Proof. This follows from Theorem 1.1 in [7] and its remark.

□

Definition 3.3. Let $X = S^1 \vee S^1 \vee \cdots \vee S^1$ and $Y = S^1 \vee S^1 \vee \cdots \vee S^1$ be identified with m and r copies of unit circle with a common point 1, respectively. Let $\pi_i : Y \to S^1$ (the ith copy of Y) be defined as follows:

$$\pi_i((\xi_1, \xi_2, ..., \xi_i, ..., \xi_r)) = \xi_i.$$

A continuous map $s : X \to Y$ is said be *standard* if $\pi_i \circ s$ on each S^1 is described as $z \to z^k$ for some integer k, where z is the identity map on the unit circle. Note that if $k = 0$, $z \to z^0$ is the map to the common point 1.

Suppose that $\kappa : \mathbb{Z}^r \to \mathbb{Z}^m$ is the homomorphism associated with a $m \times r$ matrix $(c_{i,j})$ with integer entries. Then we say that $s : X \to Y$ is the *standard map associated with* κ if $\pi_j \circ s$ on the i-th copy of S^1 can be described by $z \to z^{c_{i,j}}$.

Note that, for a standard map $s : X \to Y$, s maps the common point 1 of X to the common point 1 of Y.

It should be also noted that if $Z = S^1 \vee S^1 \vee \cdots \vee S^1$ is identified with k copies of the unit circle with a common point 1 and $s' : Y \to Z$ is a standard map, then $s' \circ s$ is a standard map from X to Z.

Lemma 3.4. *Let A be a unital simple $A\mathbb{T}$-algebra with real rank zero. Suppose that $K_1(A) = \lim_{n\to\infty}(G_n, \kappa_n) \neq \{0\}$, where G_n is direct sum of $\gamma(n)$ copies of $\mathbb{Z}$ and $\kappa_n : G_n \to G_{n+1}$ is an homomorphism. Then*

$$A = \lim_{n\to\infty}(A_n, \phi_n)$$

such that

(1) *$A_n = C(X_n) \otimes M_{k(n)} \oplus B_n$, where B_n is a finite dimensional C^*-subalgebras,*

(2) *$X_n = S^1 \vee S^1 \vee \cdots \vee S^1$, where S^1 repeats $\gamma(n)$ times,*

(3)

$$\lim_{n\to\infty} \sup_{\tau\in T(A)} \{\tau(\phi_{n,\infty}(d_n))\} = 0,$$

where d_n is the identity of $C(X_n) \otimes M_{k(n)}$,

(4) *each ϕ_n is injective.*

Moreover, if $\{X_n\}$ is fixed and a sequence of positive integers $\{a_n\}$ is given, we may require that

(5) *the partial map of ϕ_n from $C(X_n) \otimes M_{k(n)}$ to $C(X_{n+1}) \otimes M_{k(n+1)}$ has the form*

$$\phi_n(f) = \mathrm{diag}(f \circ s_n, f(\zeta(n,1)), f(\zeta(n,2)), ..., f(\zeta(n,t(n)))),$$

where s_n is a standard map associated with κ_n and $\{\zeta(n,1), \zeta(n,2), ..., \zeta(n,t(n))\}$ contains a subset that divides the j-th circle into $b(n,j)$ even arcs with $b(n,j) \geq a_n$;

(6) *each partial map $\phi_n^{(1,i)}$ of ϕ_n from $C(X_n) \otimes M_{k(n)}$ to a summand of B_{n+1} has the form*

$$\phi_n^{(1,i)}(f) = \oplus_{j=1}^{\gamma(n)} \mathrm{diag}(f(\xi(j,1)), f(\xi(j,2)), ..., f(\xi(j,k_j))),$$

where $\{\xi(j,1), ..., \xi(j,k_j)\}$ is k_j points on the j-th circle of X_n.

Proof. Note that A is an infinite dimensional simple C^*-algebra. Let B be a unital simple separable AF-algebra such that

$$(K_0(B), K_0(B)_+, [1_B]) = (K_0(A), K_0(A)_+, [1_A]).$$

By Lemma 3.1, we may assume that $B = \lim_{n\to\infty}(M_{k(n)} \oplus B_n, \phi_n)$, where B_n is a finite dimensional C^*-algebra and (1) and (2) in Lemma 3.1 hold. By passing to a subsequence, one may assume that the partial map of ϕ_n from $M_{k(n)}$ to $M_{k(n+1)}$ has multiplicity at least $m(n) \geq \gamma(n)a_n - 1$.

Put $X_n = S^1 \vee S^1 \vee \cdots \vee S^1$, where S^1 repeats $\gamma(n)$ times. Define $A_n = C(X_n) \otimes M_{k(n)} \oplus B_n$. So $K_1(A_n) = G_n = \mathbb{Z}^{\gamma(n)}$, $n = 1, 2, ...$ Define $s_n : X_n \to X_{n+1}$ to be the standard map associated with κ_n as defined in Definition 3.3. Define $\psi_n^{(1,0)}$ from $C(X_n) \otimes M_{k(n)}$ to $C(X_{n+1}) \otimes M_{k(n+1)}$ as follows:

$$\psi_n^{(1,0)}(f) = \mathrm{diag}(f \circ s_n, f(\zeta(n,1)), f(\zeta(n,2)), ..., f(\zeta(n,t(n))))$$

where $t(n) = m(n) - 1$ and $\{\zeta(n,1), \zeta(n,2), ..., \zeta(n,t(n))\}$ contains a subset that divides the j-th circle into $b(n,j)$ even arcs with $b(n,j) \geq a_n$ for each j.

By identifying $M_{k(n+1)}$ with $\mathbb{C}\otimes M_{k(n+1)}$, we also assume that $\psi_n^{(1,0)}(1_{C(X_n)\otimes M_{k(n)}}) = \phi_n^{(1,0)}(1_{M_{k(n)}})$, where $\phi_n^{(1,0)}$ is the partial map of ϕ_n from $M_{k(n)}$ to $M_{k(n+1)}$. Let $m(n,1,i)$ be the multiplicity of the partial map of ϕ_n from $M_{k(n)}$ to the i-th summand of B_{n+1}. We may assume that $m(n,1,i) \geq \gamma(n)n$. We then define

$$\psi_n^{(1,i)}(f) = \phi_n^{(1,i)}(f) = \oplus_{j=1}^{\gamma(n)} \mathrm{diag}(f(\xi(j,1)), f(\xi(j,2)), ..., f(\xi(j,k_j))),$$

where $\{\xi(j,1),...,\xi(j,k_j)\}$ is a set of k_j points on the j-th circle of X_n, from $C(X_n)\otimes M_{k(n)}$ into the i-th summand of B_n. We choose k_j so that $\sum_{j=1}^{\gamma(n)} k_j = m(n,1,i)$. Moreover, we choose $\{\xi(j,1),...,\xi(j,k_j)\}$ so that it is $2\pi/n$ dense in (the j-th) S^1. We may also assume that $\psi_n^{(1,i)}(1_{C(X_n)\otimes M_{k(n)}}) = \phi_n^{(1,0)}(1_{M_{k(n)}})$.

Now define $\psi_n : A_n \to A_{n+1}$ as follows: Define the partial map of ψ_n from $C(X_n)\otimes M_{k(n)}$ to $C(X_{n+1})\otimes M_{k(n+1)}$ to be $\psi_n^{(1,0)}$. Define the partial map of ψ_n from $C(X_n)\otimes M_{k(n)}$ to the i-th summand of B_{n+1} to be $\psi_n^{(1,i)}$. Define $\psi_n|_{B_n} = \phi_n|_{B_n}$. Note, here, we identify $M_{k(n+1)}$ with $\mathbb{C}\otimes M_{k(n+1)} \subset C(X_{n+1})\otimes M_{k(n+1)}$.

It is standard and easy to verify that $C = \lim_{n\to\infty}(A_n, \psi_n)$ is a unital simple C^*-algebra with tracial rank zero (for example, 3.7.8 and 3.7.9 of [19]

Note that

$$(K_0(A_n), K_0(A_n)_+, [1_{A_n}]) = (K_0(M_{k(n)}\oplus B_n), K_0(M_{k(n)}\oplus B_n)_+, [1_{M_{k(n)}\oplus B_n}]).$$

Moreover $(\psi_n)_{*0} = (\phi_n)_{*0}$. One also have

$$K_1(A_n) = G^{\gamma(n)}$$

and $(\psi_n)_{*1} = \kappa_n$. It follows that

$$(K_0(C), K_0(C)_+, [1_C], K_1(C)) = (K_0(A), K_0(A)_+, [1_A], K_1(A)).$$

Since C has tracial rank zero and is an AH-algebra (so it satisfies the UCT), by [21], $C \cong A$. The lemma follows from the construction. ☐

Theorem 3.5. *Let A be a unital simple $A\mathbb{T}$-algebra with real rank zero and $h : K_1(A) \to K_1(A)$ be an isomorphism. Then there exists an automorphism $\alpha : A \to A$ which satisfies the tracial cyclic Rokhlin property such that $\alpha_{*1} = h$ and $\alpha_{*0} = \mathrm{id}_{K_0(A)}$.*

Proof. If $K_1(A) = \{0\}$, it suffices to show that there exists an automorphism on A which has the Rokhlin property. But that follows from [31]. Therefore, for the rest of the proof, we may assume that $K_1(A) \neq \{0\}$.

Since $K_1(A)$ is tosion free, we may write that $K_1(A) = \lim_{n\to\infty}(G_n, \kappa_n)$, where each G_n is a direct sum of finitely many copies of $\mathbb{Z}$ and κ_n are injective.

Let $h : K_1(A) \to K_1(A)$ be the given isomorphism and let h^{-1} be the inverse. We may assume, by passing to a subsequence if necessary, that there are homomorphisms $h_n, \bar{h}_n : G_n \to G_{n+1}$ such that

$$h \circ \kappa_{n,\infty} = \kappa_{n+1,\infty} \circ h_n \quad \text{and} \quad h^{-1} \circ \kappa_{n,\infty} = \kappa_{n+1,\infty} \circ \bar{h}_n$$

on G_n. By passing to a subsequence if necessary, we may assume that,

$$h_{n+1} \circ \kappa_n = \kappa_{n+1} \circ h_n, \;\; \bar{h}_{n+1} \circ h_n = \kappa_{n+1} \circ \kappa_n \quad \text{and} \quad h_{n+1} \circ \bar{h}_n = \kappa_{n+1} \circ \kappa_n. \qquad \text{(e 3.3)}$$

Suppose that G_n is $c(n)$ copies of $\mathbb{Z}$. Homomorphisms h_n, h_n^{-1} and κ_n can be represented by matrices $(a_{i,j}^{(n)})$, $(b_{i,j}^{(n)})$ and $(c_{i,j}^{(n)})$ where $a_{i,j}^{(n)}$, $b_{i,j}^{(n)}$ and $c_{i,j}^{(n)}$ are integers.

Let $J(n) = \max\{|a_{i,j}^{(n)}| + |b_{i',j'}^{(n)}| + |c_{i'',j''}^{(n)}| : i, j, i', j', i'', j''\}$.

We identify X_n with γ_n copies of the unit circle with a common point at 1. Denote $u(n,i) \in C(X_n)$ the function on X_n which is the identity map on the ith circle and 1 everywhere else.

Define

$$\beta_n(u(n,i)) = \prod_{j=1}^{\gamma(n+1)} z(n+1,j,i),$$

where $z(n+1,j,i) = u(n+1,j)^{a^{(n)}_{i,j}}$, if $a^{(n)}_{i,j} \neq 0$, and $z(n+1,j,i) = u(n,i)(1)$ if $a^{(n)}_{i,j} = 0$, $i = 1,2,...,\gamma(n)$.

Note β_n gives a homomorphism from $C(X_n) \to C(X_{n+1})$ such that $(\beta_n)_{*1} = h_n$.

Define

$$\bar{\beta}_n(u(n,i)) = \prod_{j=1}^{\gamma(n+1)} z'(n+1,j,i),$$

where $z'(n+1,j,i) = u(n+1,j)^{b^{(n)}_{i,j}}$, if $b^{(n)}_{i,j} \neq 0$, and $z'(n+1,j,i) = u(n,i)(1)$ if $b^{(n)}_{i,j} = 0$, $i = 1,2,...,\gamma(n)$.

We now write $A = \lim_{n\to\infty}(A_n, \phi_n)$, where $A_n = C(X_n) \otimes M_{k(n)} \oplus B_n$ which satisfying (1)-(6) in Lemma 3.4 with $a_n = J(n)n$ We will also use $\beta_n : A_n \to A_{n+1}$ for the homomorphism defined by $(\beta_n)|_{C(X_n)\otimes M_{k(n)}} = \beta_n \otimes \mathrm{id}_{M_{k(n)}}$ and $(\beta_n)|_{B_n} = \phi_n|_{B_n}$.

Let $\mathcal{F}_n \subset \mathcal{F}_{n+1}$ be a sequence of finite subsets of A such that the union of those finite subsets is dense in A. Without loss of generality, we may assume that $\mathcal{F}_n \subset \phi_{n,\infty}(\mathcal{G}_n)$, where $\mathcal{G}_n$ is a finite subset of A_n.

It follows from [31] that there is an approximately inner automorphism $\sigma \in Aut(A)$ which has the tracial Rokhlin property. It follows from [25] that σ has tracial cyclic Rokhlin property.

Since σ is approximately inner, by passing to a subsequence if necessary, by Lemma 3.4, we may assume that, there is a unitary $v_n \in A_{n+1}$ such that

$$\sigma \circ \phi_{n,\infty}(f) \approx_{1/2^{n+2}} \mathrm{ad}\,\phi_{n+1,\infty}(v_n) \circ \phi_{n,\infty}(f) \quad \text{and} \tag{e 3.4}$$

$$\sigma^{-1}\phi_{n,\infty}(f) \approx_{1/2^{n+2}} \mathrm{ad}\,\phi_{n+1,\infty}(v_n^*) \circ \phi_{n,\infty}(f) \ \text{ on } \mathcal{G}_n. \tag{e 3.5}$$

Define $\sigma_n : A_{n+1} \to A_{n+1}$ by $\sigma_n(f) = \mathrm{ad}\, v_n(f)$ and $\delta_n : A_{n+1} \to A_{n+1}$ by $\delta_n(f) = \mathrm{ad}\, v_n^*(f)$ for $f \in A_n$.

Since $a_n = J(n)n$, by passing to a subsequence if necessary, we may assume that $\phi_n^{(1,1)} : C(X_n) \otimes M_{k(n)} \to C(X_{n+1}) \otimes M_{k(n+1)}$ has the following form:

$$\phi_n^{(1,1)}(f) = \mathrm{diag}(f \circ s_n, f(\zeta(n,1)), f(\zeta(n,2),..., f(\zeta(n,t(n))),$$

where s_n is a standard map (see Definition 3.3) and $\{\zeta(n,1)), \zeta(n,2),...,\zeta(n,t(n)\}$ contains subsets $\{\xi_1,...,\xi_N\}$, $\{\xi_{N+1},....\xi_{2N}\}$,...,$\{\xi_{(\gamma(n)-1)N+1},...,\xi_{\gamma(n)N}\}$ such that $\{\xi_{(j-1)N+1},...,\xi_{jN}\}$ evenly divide the j-th copy of S^1 and $N = \Lambda(n, \mathcal{G}_n)$.

Define a monomorphism $\omega_n : C(X_n) \otimes M_{k(n)} \to A_{n+2}$ as follows: Define

$$\omega_n^{(1,1)}(f) = \beta_{n+1}(g_1) + \phi_{n+1}(g_2),$$

where $g_1 = \sigma_n(\mathrm{diag}(f \circ s_n, 0 \cdots 0))$ and $g_2 = \sigma_n(0, f(\zeta(n,1)), f(\zeta(n,2)),..., f(\zeta(n,t(n)))$. Define $\omega_n^{(1,i)} = \phi_{n+1}^{(1,i)} \circ \phi_n|_{C(X_n)\otimes M_{k(n)}}$. Let $E_{n+2} = \phi_{n,n+2}(d_n)$, where d_n is the identity of $C(X_n) \otimes M_{k(n)}$. Denote by

$$E'_{n+1} = \mathrm{diag}(1, \overbrace{0,...,0}^{t(n)}) \in C(X_{n+1}) \otimes M_{k(n+1)}.$$

Define a monomorphism $\bar{\omega}_n : C(X_n) \otimes M_{k(n)} \to E_{n+2}A_{n+2}E_{n+2}$ as follows:

Define
$$\bar{\omega}_n^{(1,1)}(f) = \bar{\beta}_{n+1}(g_1') + \phi_{n+1}(g_2'),$$
where $g_1' = \delta_n(\mathrm{diag}(f \circ s_n, 0 \cdots 0))$ and $g_2' = \delta_n(0, f(\zeta(n,1)), f(\zeta(n,2), ..., f(\zeta(n,t(n))))$. Define $\bar{\omega}_n^{(1,i)} = \phi_{n+1}^{(1,i)} \circ \phi_n|_{C(X_n)\otimes M_{k(n)}}$.

For any unitary $u \in A_{4n+3}$, define $\Phi : C(X_{4n+1}) \otimes M_{k(4n+1)} \to A_{4n+5}$ by
$$\Phi(f) = \bar{\beta}_{4n+4} \circ \delta_{4n+3} \circ \phi_{4n+3} \circ \mathrm{ad}\, u \circ \beta_{4n+2}(g_1).$$
Since
$$(\bar{\beta}_{4n+4} \circ \delta_{4n+3} \circ \phi_{4n+3} \circ \mathrm{ad}\, u \circ \beta_{4n+2} \circ \phi_{4n+1})_{*1} = (\phi_{4n+1,4n+5})_{*1},$$
$$(\Phi)_{*1} = (\phi_{4n+2,4n+5} \circ E_{4n+1}'\phi_{4n+1}E_{4n+1}')_{*1}.$$

By choice of $\Lambda(n, \mathcal{G}_n)$, by (e 3.3) and by applying Lemma 3.2, we obtain unitaries $u_k \in E_kA_kE_k$ such that (with $u_1 = d_1$)
$$\mathrm{ad}\, u_{4(n+1)+1} \circ \bar{\omega}_{4n+3} \circ \mathrm{ad}\, u_{4n+3} \circ \omega_{4n+1}(f) \approx_{1/2^n} \phi_{4n+1,4(n+1)+1}(f) \text{ on } d_{4n+1}\mathcal{G}_{4n+1} \qquad \text{(e 3.6)}$$
as well as
$$\mathrm{ad}\, u_{4(n+1)+3} \circ \omega_{4(n+1)+3} \circ \mathrm{ad}\, u_{4n+3} \circ \bar{\omega}_{4n+3}(f) \approx_{1/2^n} \phi_{4n+3,4(n+1)+3}(f) \text{ on } d_{4n+3}\mathcal{G}_{4n+3} \qquad \text{(e 3.7)}$$
$(n = 0, 1, 2, ...)$

Define $\omega_{4n+1}' = \mathrm{ad}\, u_{4n+3} \circ \omega_{4n+1}$ and $\overline{\varpi}'{}_{4n+3} = \mathrm{ad}\, u_{4n+5} \circ \bar{\omega}_{4n+3}$.

Now define $\alpha_n : A_{4n+1} \to A_{4n+3}$ by
$$(\alpha_n)|_{C(X_{4n+1})\otimes M_{k(4n+1)}} = \omega_{4n+1}' \quad \text{and} \quad (\alpha_n)|_{B_{4n+1}} = (\phi_{4n+2} \circ \sigma_{4n+1} \circ \phi_{4n+1})|_{B_{4n+1}}$$
and define $\bar{\alpha}_n : A_{4n+3} \to A_{4n+5}$ by
$$(\bar{\alpha}_n)|_{C(X_{4n+3})\otimes M_{k(4n+3)}} = \overline{\omega'}_{4n+3} \quad \text{and} \quad (\bar{\alpha}_n)|_{B_{4n+3}} = (\phi_{4n+4} \circ \delta_{4n+3} \circ \phi_{4n+3})|_{B_{4n+3}}$$

From (e 3.4) and (e 3.5), we have that
$$\bar{\alpha}_{n+1} \circ \alpha_n(f) \approx_{1/2^{n+1}} \phi_{4n+1,4(n+1)+1}(f) \text{ on } f \in (1 - d_{4n+1})\mathcal{G}_{4n+1} \qquad \text{(e 3.8)}$$

$$\alpha_{n+1} \circ \bar{\alpha}_n(f) \approx_{1/2^{n+1}} \phi_{4n+3,4(n+1)+3}(f) \text{ on } f \in (1 - d_{4n+1})\mathcal{G}_{4n+1} \qquad \text{(e 3.9)}$$

It follows from (e 3.6), (e 3.7), (e 3.8) and (e 3.9), one has the following approximately intertwining:
$$\begin{tikzcd}
A_1 \arrow[rr, "\phi_{1,5}"] \arrow[dr, "\alpha_1"] & & A_5 \arrow[rr, "\phi_{5,9}"] \arrow[dr, "\alpha_2"] & & A_9 \arrow[rr, "\phi_{9,13}"] \arrow[dr, "\alpha_3"] & & A_{17}\cdots \\
& A_3 \arrow[rr, "\phi_{3,7}"] \arrow[ur, "\bar{\alpha}_1"] & & A_7 \arrow[rr, "\phi_{7,11}"] \arrow[ur, "\bar{\alpha}_2"] & & A_{11} \arrow[rr, "\phi_{11,15}"] \arrow[ur, "\bar{\alpha}_3"] & & \cdots
\end{tikzcd}$$

It follows from the Elliott approximately intertwining argument that $\{\alpha_n\}$ and $\{\bar{\alpha}_n\}$ define two automorphisms α and α^{-1}.

Moreover, one checks that

$$\alpha_{*0} = \mathrm{id}_{K_0(A)} \quad \text{and} \quad \alpha_{*1} = h.$$

It remains to show that α has tracial cyclic Rokhlin property.

Fix $\varepsilon > 0$, $m > 0$ and a finite subset $\mathcal{F} \subset A$. Since σ has tracial cyclic Rokhlin property, there are mutually orthogonal projections $e_1, e_2, ..., e_m$ such that (with $e_{m+1} = e_1$)

$$\|e_i a - a e_i\| < \varepsilon/8, \ \ i = 1, 2, ..., m, a \in \mathcal{F} \tag{e 3.10}$$

$$\|\sigma(e_i) - e_{i+1}\| < \varepsilon/8, \ \ i = 1, 2, ..., m \tag{e 3.11}$$

and

$$\tau(1 - \sum_{i=1}^{m} e_i) < \varepsilon/8 \tag{e 3.12}$$

for all $\tau \in T(A)$.

Since σ is approximately inner, there is a unitary $v \in A$ such that

$$\|v^* e_i v - e_{i+1}\| < \varepsilon/8, \ \ i = 1, 2, ..., m.$$

Without loss of generality, we may assume that there is a finite subset $\mathcal{G} \subset A_{4n+1}$ such that $\mathcal{F} \subset \phi_{4n+1,\infty}(\mathcal{G})$ for some sufficiently large n.

It follows from the standard argument (See [34, Proposition L.2.2] for example.) that for any $\delta > 0$, (for sufficiently large n) there are projections $q_i \in A_{4n+1}$ such that

$$\|e_i - \phi_{4n+1,\infty}(q_i)\| < \min\{\delta, \varepsilon/8\}, \ \ i = 1, 2, ..., m. \tag{e 3.13}$$

Since $\phi_{n,\infty}$ is assumed to be injective, with sufficiently small δ, from [19, Lemma 2.5.6] we may assume that $q_1, q_2, ..., q_m$ are mutually orthogonal. Let $\bar{q}_i = \phi_{4n+1,\infty}(q_i)$. We may also assume that there is a unitary $v' \in A_{4n+1}$ such that $\phi_{4n+1,\infty}(v') = v$.

With even larger n, we may assume that, without loss of generality,

$$\alpha^{-i}(\mathcal{F}),\ \alpha^{-i}(\bar{q}_i) \in \phi_{4n+1,\infty}(A_{4n+1}), \ \ i = 1, 2, ..., m. \tag{e 3.14}$$

Note that $(1 - d_{4n+1})$ commutes with every element in A_{4n+1}. Put $p_i = \bar{q}_i \alpha^{i-1}(\phi_{4n+1,\infty}(1 - d_{4n+1}))$, $i = 1, 2, ..., m$. Since, by (e 3.14),

$$\alpha^{i-1} \circ \phi_{4n+1,\infty}(1 - d_{4n+1})\bar{q}_i = \alpha^{i-1}(\phi_{4n+1,\infty}(1 - d_{4n+1})\alpha^{-i+1}(\bar{q}_i))$$
$$= \alpha^{i-1}(\alpha^{-i+1}(\bar{q}_i)\phi_{4n+1,\infty}(1 - d_{4n+1})) = \bar{q}_i \alpha^{i-1} \circ \phi_{4n+1,\infty}(1 - d_{4n+1}),$$

$\{p_1, p_2, ..., p_m\}$ are mutually orthogonal projections.

By (e 3.14), if $a \in \mathcal{F}$,

$$p_i a = \bar{q}_i \alpha^{i-1}(\phi_{4n+1,\infty}(1 - d_{4n+1})\alpha^{-i+1}(a)) = \bar{q}_i a \alpha^{i-1}(\phi_{4n+1,\infty}(1 - d_{4n+1})).$$

Therefore, by (e 3.13) and (e 3.10), we have

(1) $\|p_i a - a p_i\| = \|\bar{q}_i a \alpha^{i-1}((\phi_{4n+1,\infty}(1 - d_{4n+1}))) - a p_i\| < \varepsilon/8 + \varepsilon/8 + \varepsilon/8 < \varepsilon, \ \ i = 1, 2, ..., m.$

Since

$$\alpha(p_i)=\alpha(\bar{q}_i\alpha^{i-1}(\phi_{4n+1,\infty}(1-d_{4n+1})))=\alpha(\bar{q}_i)\alpha^i(\phi_{4n+1,\infty}(1-d_{4n+1}))),$$

by (e 3.10) and (e 3.11),
(2) $\|\alpha(p_i)-p_{i+1}\|=\|(\alpha(\bar{q}_i)-\bar{q}_{i+1})\alpha^i(\phi_{4n+1,\infty}(1-d_{4n+1}))\|<\varepsilon$, $i=1,2,...,m$.
Note, since

$$\lim_{n\to\infty}\sup\{\tau(\phi_{n,\infty}(d_n)):\tau\in T(A)\}=0,$$

we may assume that

$$\tau(\phi_{4n+1,\infty}(d_{4n+1}))<\varepsilon/4m$$

for all $\tau\in T(A)$.

Therefore, using the fact that
$\alpha_{*0}=\mathrm{id}_{K_0(A)}$,

$$\tau(p_i)=\tau(\bar{q}_i\alpha^{i-1}(\phi_{4n+1,\infty}(1-d_{4n+1})))>\tau(\bar{q}_i)(1-\varepsilon/4m),\ \ i=1,2,...,m$$

Note also, by (e 3.13) and with $\delta<1$,

$$\tau(\bar{q}_i)=\tau(e_i),\ \ i=1,2,...,m.$$

Applying (e 3.12), we also have
(3)

$$\begin{aligned}\tau(1-\sum_{i=1}^m p_i) &< 1-(1-\frac{\varepsilon}{4m})\sum_{i=1}^m\tau(\bar{q}_i)\\ &= 1-(1-\frac{\varepsilon}{4m})\sum_{i=1}^m\tau(e_i)\\ &= (1-\sum_{i=1}^m\tau(e_i))+\frac{\varepsilon}{4m}\sum_{i=1}^m\tau(e_i)\\ &< \frac{\varepsilon}{8}+\frac{\varepsilon}{4}\tau(e_1)<\frac{\varepsilon}{8}+\frac{\varepsilon}{4}<\varepsilon\end{aligned}$$

for all $\tau\in T(A)$.

□

Corollary 3.6. *Let A be a unital simple $A\mathbb{T}$-algebra with real rank zero and let $h:K_1(A)\to K_1(A)$ be an isomorphism. Suppose that α is the automorphism given by Theorem 3.5. Then the crossed product $A\rtimes_\alpha\mathbb{Z}$ has tracial rank zero and can be written as an AH-algebra.*

Proof. Since $\alpha_{*0}=\mathrm{id}_{K_0(A)}$ and has the tracial cyclic Rokhlin property, by [24] $A\rtimes_\alpha\mathbb{Z}$ has tracial rank zero. It follows from the classification theorem [21] that $A\rtimes_\alpha\mathbb{Z}$ is in fact simple AH-algebra. □

4 Minimal dynamical systems and non-commutative dynamical systems

It seems that a more general construction based on Theorem 3.5 can be made so that Theorem 3.5 can covers all simple AH-algebras with real rank zero and stable rank one. However, in this section, we present a different type of construction. If one believes that automorphisms on C^*-algebras with Rokhlin property is closely related to minimal homeomorphisms on compact metric spaces, then the construction that we presented here sheds some light on these relation. Given a connected finite CW complex X, there is a standard way to construct a unital simple C^*-algebra with similar topological information:

Proposition 4.1. *Let X be a connected finite CW-complex space and let D be a dimension group which is a countable dense subgroup of $\mathbb{Q}$. Let $\rho_X : K_0(C(X)) \to \mathbb{Z}$ be the dimension map. Then there exists a unital separable AH-algebra A_X with tracial rank zero and with a unique tracial state τ such that*

$$(K_0(A_X), K_0(A_X)_+) = D \oplus \mathrm{ker}\rho_X,\ \mathrm{ker}\rho_{A_X} = \mathrm{ker}\rho_X \quad \text{and} \quad K_1(A_X) = K_1(C(X)).$$

Moreover, there is a unital embedding $j : C(X) \to A_X$ such that

$$j_{*0}|_{\mathrm{ker}\rho_X} = \mathrm{id}_{\mathrm{ker}\rho_X} \quad \text{and} \quad j_{*1} = \mathrm{id}_{K_1(C(X))}.$$

(See [20] for the construction).

Suppose that there is a minimal homeomorphism $\psi : X \to X$. One can then construct an automorphsim α on A_X associated with ψ.

Theorem 4.2. *Let X be a connected finite CW-complex space, and let $\psi\colon X \to X$ be a minimal homeomorphism. Denote by $\psi^\natural : C(X) \to C(X)$ the automorphism defined by $\psi^\natural(f) = f \circ \psi$. Then there is an automorphism $\alpha : A_X \to A_X$ which satisfies the tracial cyclic Rokhlin property such that*

$$\alpha_{*0}|_{\mathrm{ker}\rho_X} = (\psi^\natural)_{*0}|_{\mathrm{ker}\rho_X} \quad \text{and} \quad \alpha_{*1} = (\psi^\natural)_{*1}.$$

Theorem 4.2 is a corollary of a more general Theorem 4.10 below.

Lemma 4.3. *Let X be a compact connected metric space and let $\psi : X \to X$ be a minimal homeomorphism with a ψ-invariant Borel probability measure μ. Let S be an infinite subset of positive integers. Then, for any $\varepsilon > 0$, there is an integer $n \in S$ and there is a finite ε-dense set $\{x_1, x_2, ..., x_n\} \subset X$ such that for any closed subset $F \subset X$,*

$$\mu_1(F) \le \mu_2(F_\varepsilon) \quad \text{and} \quad \mu_2(F) \le \mu_1(F_\varepsilon), \tag{e 4.15}$$

where μ_1 is the measure concentrated on $\{x_1, x_2, ..., x_n\}$ with $\mu_1(\{x_j\}) = \frac{1}{n}$ and μ_2 is the measure concentrated on $\{\psi(x_1), \psi(x_2), ..., \psi(x_n)\}$ with $\mu_2(\{\psi(x_j)\}) = \frac{1}{n}$.

Proof. First we note that any open neighborhood $O(x)$ of any $x \in X$ must have positive μ measure. Otherwise, assume that $\mu(O(x)) = 0$. Put

$$O = \cup_{n\in\mathbb{Z}}\psi^n(O(x)).$$

Then O is open and $\psi(O) = O$. It follows that $F = X \setminus O$ is invariant under ψ. The minimality of ψ implies that $F = \emptyset$. This implies that $O = X$. But $\mu(O) = 0$. A contradiction.

Fix $\delta > 0$. Since X is compact, by what we have just proved, it is easy to find finitely many disjoint Borel subsets $G_1, G_2, ..., G_m$ such that $\mu(G_i) > 0$ and $\mathrm{diam}(G_i) < \delta$ for $i = 1, 2, ..., m$, and

$$\cup_{i=1}^m G_i = X.$$

Let $a = \min\{\mu(G_i) : 1 \le i \le m\}$. Choose $n \in S$ so that

$$\frac{1}{n} < \frac{a}{2m^{m+1}}.$$

There is an integer $0 < k_i < n$ such that

$$\frac{k_i}{n} \le \mu(G_i) < \frac{k_i+1}{n}, i = 1, 2, ..., m.$$

By minimality of ψ, no single point has positive μ-measure. Thus each G_i contains infinitely many points. Choose k_i points in G_i, $i = 1, 2, ..., m-1$. Then

$$\sum_{i=1}^m k_i \ge n - m.$$

Choose $k_m' = k_m + (n - \sum_{i=1}^m k_i)$ points in G_m. Note that

$$\frac{k_m'}{n} - \mu(G_m) \le \frac{m}{n}. \tag{e 4.16}$$

We obtain n points $x_1, x_2, ..., x_n$ in X. Now define $\mu_{\delta,1}$ to be the measure concentrated on $\{x_1, x_2, ..., x_n\}$ with $\mu_{\delta,1}(\{x_j\}) = \frac{1}{n}$ and define $\mu_{\delta,2}$ to be the measure concentrated on $\{\psi(x_1), \psi(x_2), ..., \psi(x_n)\}$ with $\mu_{\delta,2}(\{\psi(x_j)\} = \frac{1}{n}$. It is clear that when $\delta \to 0$, $m \to \infty$. Fix $\xi(\varepsilon, j) \in G_j$. Note that for each $f \in C(X)$,

$$\sum_{j=1}^m f(\xi(\delta, j))\mu(G_j) \to \int_X f d\mu$$

as $\delta \to 0$. It follows that

$$\int_X f d\mu_{\delta,1} \to \int_X f d\mu$$

for all $f \in C(X)$ as $\delta \to 0$. We also have

$$\int_X f\mu_{\delta,2} = \int_X f \circ \psi d\mu_{\delta,1} \to \int_X f \circ \psi d\mu = \int_X f d\mu$$

for all $f \in C(X)$.

Now let $\varepsilon > 0$ and $y_1, y_2, ..., y_K$ be a subset of X such that

$$\cup_{j=1}^K O(y_j, \varepsilon/4) = X.$$

Let $\mathcal{O}$ be the finite collection of all possible union of $O(y_j, \varepsilon/4)$. Let $\mathcal{O}'$ be the finite collection of all possible union of $O(y_j, \varepsilon/2)$. If $G \subset \mathcal{O}$ we denote by G' the corresponding union of $O(y_j, \varepsilon/2)$'s.

Let

$$b = \min\{\mu(G') - \mu(G) : G \subset \mathcal{O} \quad \text{and} \quad G \neq X\}.$$

The connectedness of X together the first paragraph of this proof imply that $b > 0$.

Choose δ sufficiently small so that, for all $G \subset \mathcal{O}$

$$\mu_{\delta,i}(G) - \frac{b}{4} \leq \mu(G)) \leq \mu_{\delta,i}(G) + \frac{b}{4}$$

and, for any $G' \in \mathcal{O}'$,

$$\mu_{\delta,i}(G') - \frac{b}{4} \leq \mu(G') \leq \mu_{\delta,i}(G') + \frac{b}{4}$$

$i = 1, 2$. Thus, if $G \neq X$,

$$\mu_{\delta,i}(G) + \frac{b}{4} \leq \mu(G) + \frac{b}{4} \leq \mu(G') - \frac{b}{4} \leq \mu_{\delta,i}(G'), i = 1, 2.$$

Now let F be a closed subset of X. If $F_\varepsilon = X$, then (e 4.15) follows. Now suppose that $F_\varepsilon \neq X$. Let

$$G = \cup_{F \cap O(y_j, \varepsilon/4) \neq \emptyset} O(y_j, \varepsilon/4) \subset \mathcal{O}$$

Then $G \neq X$. It follows that

$$\mu_{\delta,1}(F) \leq \mu(G) \leq \mu_{\delta,2}(G) + \frac{b}{4} \leq \mu_{\delta,2}(G') \leq \mu_{\delta,2}(F_\varepsilon).$$

Similarly

$$\mu_{\delta,2}(F) \leq \mu_{\delta,1}(F_\varepsilon).$$

□

Lemma 4.4. *Let X be a compact connected metric space (with infinitely many points) and let $\psi : X \to X$ be a minimal homeomorphism. Let S be an infinite subset of $\mathbf{N}$. Then, for any $\varepsilon > 0$, there is $n \in S$ and there is a finite ε-dense set $\{x_1, x_2, ..., x_n\} \subset X$ such that there is a permutation $s : (1, 2, ..., n) \to (1, 2, ..., n)$ such that*

$$\text{dist}(x_j, \psi(x_{s(j)})) < \varepsilon, j = 1, 2, ..., n.$$

Proof. Let μ be a ψ-invariant Borel probability measure. Let $\{x_1, x_2, ..., x_n\}$ be a finite subset satisfying the conclusion of Lemma 4.3.

For any subset S_1 of l elements in $\{x_1, x_2, ..., x_n\}$, choose a closed subset $F \cap \{x_1, x_2, ..., x_n\} = S_1$ and

$$F \subset \{x \in X : \text{dist}(x, \{x_1, x_2, ..., x_n\}) < \varepsilon/4\}.$$

Then

$$\mu_1(F) \leq \mu_2(F_{\varepsilon/4}).$$

Thus $F_{\varepsilon/4}$ contains at least l elements in $\{\psi(x_1), \psi(x_2), ..., \psi(x_n)\}$. Therefore there is a subset S_2 of l elements in $\{\psi(x_1), \psi(x_2), ..., \psi(x_n)\}$ such that, for any element $\xi \in S_1$ there is $\xi' \in S_2$ such that

$$\text{dist}(\xi, \xi') < \varepsilon.$$

By the "Marriage Lemma" (see [9]), there is a permutation s which meets the requirement.

□

Lemma 4.5. *Let X be a compact connected metric space (with infinitely many points) with a metric d, and $\psi\colon X \to X$ be a minimal homeomorphism. For any $\epsilon > 0$ and a finite set $\mathcal{F} \subset C(X)$ there exists $\delta > 0$ such that if $\{x_i\}_{i=1}^n$ are points in X which satisfy the property that there is a permutation $s\colon (1, 2, \ldots, n) \to (1, 2, \ldots, n)$ such that*

$$\mathrm{dist}(x_j, \psi(x_{s(j)})) < \delta,$$

then there is a permutation $n \times n$ matirx U such that

$$\|\mathrm{diag}(f(x_1), \ldots, f(x_n)) - U\mathrm{diag}(f(\psi(x_1)), \ldots, f(\psi(x_n))U^*\| < \epsilon, \quad \forall f \in \mathcal{F}.$$

Proof. Let $\varepsilon > 0$ and $\mathcal{F} \subset C(X)$ be a finite set. Then there exists $\delta > 0$ such that if $d(x, x') < \delta$ then $|f(x) - f(x')| < \varepsilon$ for any $f \in \mathcal{F}$.

Let $\{x_1, x_2, \ldots, x_n\}$ be points in X satisfy that there is a permutation $s\colon (1, 2, \ldots, n) \to (1, 2, \ldots, n)$ such that

$$\mathrm{dist}(x_j, \psi(x_{s(j)})) < \delta.$$

Take a permutation matrix U correspondent to s, then

$$\begin{aligned}
&\|\mathrm{diag}(f(x_1), \ldots, f(x_n)) - U\mathrm{diag}(f(\psi(x_1)), \ldots, f(\psi(x_n))U^*\| \\
&= \|\mathrm{diag}(f(x_1), \ldots, f(x_n)) - \mathrm{diag}(f(\psi(x_{s(1)})), \ldots, f(\psi(x_{s(n)}))\| \\
&< \varepsilon
\end{aligned}$$

□

Lemma 4.6. *Let X be a compact connected metric space with a minimal homeomorphism $\psi : X \to X$. Let $\{f_n\}$ be a dense sequence in $C(X)$, let $\{\varepsilon_n\}$ be a sequence of positive numbers and let $\{k(n)\}$ be a sequence of positive integers. Then there exists a sequence of positive numbers $\{\delta_n\}$ which has the following property: If $\{x(1, n), ..., x(l(n), n)\}$ are points in X which satisfy the following property: there is a permutation $s_n : (1, 2, ..., l(n)) \to (1, 2, ..., l(n))$ such that*

$$\mathrm{dist}(x(j, n), \psi(x(s_n(j), n))) < \delta_n$$

then there is a sequence of permutation $l(n) \times l(n)$ matrices U_n such that

$$W_n^*\mathrm{diag}(f(x(1, n)), f(x(2, n)), ..., f(x(l(n), n)))W_n$$

$$\approx_{\varepsilon_n} \mathrm{diag}(f(\psi(x(1, n))), f(\psi(x(2, n))), ..., f(\psi(x(l(n), n))))$$

for all $f \in \{(a_{i,j}) \in C(X) \otimes M_{k(n)} : a_{i,j} \in \{f_1, f_2, ..., f_n\} \cup \mathbb{C}\}$, where $W_n = 1_{k(n)} \otimes U_n$ is a matrix in $M_{k(n)l(n)}$.

Proof. This follows from Lemma 4.5 immdiately. □

We shall construct a model automorphism with tracial Rokhlin property on some AH algebras which contains the class of $A\mathbb{T}^n$-algebras and AI-algebras.

Let X be a compact metric space with infinitely many points. Let (X, ψ) be a minimal dynamical system. If (X, ψ) is uniquely ergodic, then (X, ψ) has mean dimension zero. If X has finite covering dimension, then (X, ψ) always has mean dimension zero ([26]).

Lemma 4.7. *Let X be a compact metric space and let $\psi : X \to X$ be a minimal homeomorphism. Suppose that (X, ψ) has mean dimension zero. Then, for any integer $N \geq 1$, any $\eta > 0$ and any $\delta > 0$, there are mutually disjoint open subsets $G_1, G_2, ..., G_L$ such that*

(1) $\mathrm{diam}(G_i) < \eta$, $i = 1, 2, ..., L$,

(2) $\psi^i(\cup_{s=1}^L G_s) \cap \psi^j(\cup_{s=1}^L G_s)) = \emptyset$, *if* $i \neq j$ *and* $i, j = 0, 1, 2, ..., N-1$,

(3) $\mu(\cup_{j=0}^{N-1} \psi^j(\cup_{i=1}^L G_i)) > 1 - \delta$ *for all* $\mu \in T_\psi$ *and*

(4) $\mu(\partial(G_i)) = 0$ *for* $i = 1, 2, \ldots, L$ *and all* $\mu \in T_\psi$, *where* T_ψ *is the set of all* ψ*-invariant Borel probability measures*

This basically follows from Lemma 3.4 and Lemma 3.5 of [23].

Lemma 4.8. *Let X be a compact metric space and let $\varepsilon > 0$. Suppose that $\mathcal{G} \subset C(X)$ satisfies the following: if* $\mathrm{dist}(x, x') < \delta$, *then*

$$|f(x) - f(x')| < \varepsilon/m^2$$

for all $f \in \mathcal{G}$. Denote by $\mathcal{F} = \{(a_{i,j}) \in C(X) \otimes M_m : a_{i,j} \in \mathcal{G}\}$. Let $\phi : C(X) \otimes M_m \to C(X) \otimes M_{m(k+l)}$ be defined by

$$\phi(f) = \mathrm{diag}(\overbrace{f, f, ..., f}^{l}, f(\xi_1), f(\xi_2), ..., f(\xi_k))$$

for all $f \in C(X) \otimes M_m$. Then if $\mathrm{dist}(x, x') < \delta$,

$$\|\phi(f)(x) - \phi(f)(x')\| < \varepsilon$$

for all $f \in \mathcal{F}$, where the norm is operator norm on $M_{m(k+l)}$.

Definition 4.9. Let X be a compact connected metric space with infinitely many points. Let $d_X : K_0(C(X)) \to \mathbb{Z}$ be the dimension map. Write $K_0(C(X)) = \mathbb{Z} \oplus G_{00}$ and $K_1(C(X)) = G_1$, where $G_{00} = \mathrm{ker} d_X$. Fix two sequences $\{a_n\}$ and $\{b_n\}$ and a sequence $\{k_n\}$ of positive integers (with $k_n \to \infty$ as $n \to \infty$). We assume that $k_1 = 1$ and $k_{n+1} = b_n k_n$, where b_n a sequence of positive numbers such that $b_n \geq 2$. Let $H_{00} = \lim_{n\to\infty}(G_{00}, \kappa_{n,n+1}^{(00)})$, where $\kappa_{n,n+1}^{(00)} : G_{00} \to G_{00}$ is defined by $\kappa_{n,n+1}^{(00)}(g) = a_n g$ for all $g \in G_{00}$. Let D be the dimension group defined by $D = \lim_{n\to\infty}(\mathbb{Z}, \kappa_{n,n+1}^{(d)})$, where $\kappa_{n,n+1}^{(d)}(g) = b_n g$ for all $g \in \mathbb{Z}$. Define $H_0 = D \oplus H_{00} = \lim_{n,\infty}(\mathbb{Z} \oplus G_{00}, \kappa_{n,n+1}^{(0)})$, where $\kappa_{n,n+1}^{(0)} = \kappa_{n,n+1}^{(d)} \oplus \kappa_{n,n+1}^{(00)}$. Define $H_1 = \lim_{n\to\infty}(G_1, \kappa_{n,n+1}^{(0)})$, where $\kappa_{n,n+1}^{(1)}(g) = a_n g$ for all $g \in G_1$. By passing to a subsequence, without changing H_{00}, D or H, one may assume that $b_n = a_n + l(n)$ and

$$\lim_{n\to\infty} \frac{a_n}{b_n} = 0.$$

There is a unital simple AH-algebra $A = A(X, \{a_n\}, \{b_n\})$ such that

$$(K_0(A), K_0(A)_0, [1_A], K_1(A)) = (D \oplus H_{00}, (D \oplus H_{00})_+, 1, H_1),$$

where

$$(D \oplus H_{00})_+ = \{(d, x) : d > 0, d \in D, x \in H_{00}\} \cup \{(0, 0)\}.$$

(see, for example, [20]). In this case, D is identified with a countable dense subgroup of $\mathbb{R}$. Note that A has a unique tracial state τ such that $\rho_A(K_0(A)) = D$. By the classification theorem ([21]), there is only one such separable simple amenable C^*-algebra with tracial rank zero satisfying the UCT.

Suppose that $\gamma_0 : K_0(C(X)) \to K_0(C(X))$ is an order isomorphism and $\gamma_1 : K_1(C(X)) \to K_1(C(X))$ is an isomorphism. Then γ_0 induces an order isomorphism $I(\gamma_0) : D \oplus H_{00} \to D \oplus H_{00}$ such that $I(\gamma_0)((d, 0)) = (d, 0)$ and if $\kappa^{(0)}_{n,\infty} : G_{00} \to H_{00}$ is the homomorphism induced by the inductive system, $I(\gamma_0) \circ \kappa^{(0)}_{n,\infty}(g) = \kappa^{(0)}_{n,\infty}(\gamma_0(g))$ for all g in the n-th $\mathbb{Z} \oplus G_{00}$. Similarly $I(\gamma_1) \circ \kappa^{(1)}_{n,\infty}(g) = \kappa_{n,\infty} \circ \gamma_1(g)$ for all g in the n-th G_1.

Theorem 4.10. *Let X be a connected compact metric space with $K_0(C(X)) = \mathbb{Z} \oplus G_{00}$ and $K_1(C(X)) = G_1$. Fix $\{a_n\}$ and $\{b_n\}$ two sequences of positive numbers. Let $A = A(X, \{a_n\}, \{b_n\})$. Suppose that X has a minimal homeomorphism ψ such that (X, ψ) has mean dimension zero. Then there exists an automorphism $\alpha : A \to A$ with the tracial cyclic Rokhlin property such that*

$$\alpha_{*i} = I(\psi^{\natural}_{*i}) \text{ on } K_i(A),$$

$i = 0, 1$.

Proof. Let $k(1) = 1$, $k(n + 1) = b_n k(n)$, $n = 1, 2, ...,$. Put $l_n = b_n - a_n$. We may assume, as mentioned in Definition 4.9, that that $\lim_{n\to\infty} \frac{a_n}{b_n} = 0$.

Choose $\varepsilon_n = \frac{1}{2^n}$. Fix a dense subset $\{f_1, f_2, ..., f_n, ...\}$. Let $\{\delta_n\}$ be as in Lemma 4.6 associated with $\{k(n)\}$, $\{f_1, f_2, ..., f_n, ...\}$ and $\{\varepsilon_n\}$. Let $F_n = \{x(1, n), x(2, n), ..., x(l(n), n)\}$ be a finite subset (with $l(n)$ elements) of X such that F_n is $1/2^n$ -dense in X and there is a permutation $s_n : (1, 2, ..., l(n)) \to (1, 2, ..., l(n))$ such that

$$\mathrm{dist}(x(j, n), \psi(x_{s_n(x(j,n))})) < \min\{\delta_n, \varepsilon_n\}\ j = 1, 2, ..., l(n).$$

Such F_n is given by Lemma 4.4.

Define $\phi_n : C(X) \otimes M_{k(n)} \to C(X) \otimes M_{k(n+1)}$ by

$$\phi_n(f) = \mathrm{diag}(\overbrace{f, f, ..., f}^{a_n}, f(x(1, n)), f(x(2, n)), ..., f(x(l(n), n))) \qquad \text{(e 4.17)}$$

for $f \in C(X) \otimes M_{k(n)}$. Then $A \cong \lim_{n\to\infty}(C(X) \otimes M_{k(n)}, \phi_n)$.

Note that A has a unique tracial state (see Definition 4.9). Denote it by τ.

Define $\psi^{\natural} : C(X) \to C(X)$ by $\psi^{\natural}(f)(x) = f(\psi(x))$ for $x \in X$. Put $A_n = C(X) \otimes M_{k(n)}$ for each $n \in \mathbf{N}$ and define $\alpha_n = \psi^{\natural} \otimes id_{k_n} : A_n \to A_n$. Then $\alpha_n \in \mathrm{Aut}(A_n)$ for each $n \in \mathbf{N}$. We set

$$\mathcal{F}_n = \{(a_{i,j}) \in M_{k(n)} : a_{i,j} \in \{f_1, f_2, ..., f_n\} \cup \mathbb{C}\}.$$

From Lemma 4.6 we may assume that there is a permutation matrix $U'_{n+1} \in M_{l(n)}$ such that

$$||\phi_n \circ \alpha_n(f) - \mathrm{ad}(U_{n+1}) \circ \alpha_{n+1} \circ \phi_n(f)|| < \varepsilon_n \qquad \text{(e 4.18)}$$

for each $n \geq 1$ and $f \in \mathcal{F}_n \cup \cup_{j=1}^{n-1} \phi_j(\mathcal{F}_j)$, where $U_{n+1} = \mathrm{diag}(1_{a_n k(n)}, 1_{k(n)} \otimes U'_{n+1}) \in M_{k(n+1)}$. Thus we obtain a permutation matrix $V'_n \in M_{l(n+1)}$ such that the following approximately intertwining:

$$\begin{tikzcd}
A_1 \arrow[r, "\phi_1"] \arrow[d, "\alpha_1"'] & A_2 \arrow[r, "\phi_2"] \arrow[d, "\mathrm{ad}V_2\circ\alpha_2"'] & A_3 \arrow[r, "\phi_3"] \arrow[d, "\mathrm{ad}V_3\circ\alpha_3"'] & \cdots \\
A_1 \arrow[r, "\phi_1"] & A_2 \arrow[r, "\phi_2"] & A_3 \arrow[r, "\phi_3"] & \cdots,
\end{tikzcd}$$

where V_n is a sequence of permutation matrixes in M_{k_n} such that

$$||\phi_n \circ \mathrm{ad}(V_n) \circ \alpha_n(f) - \mathrm{ad}(V_{n+1}) \circ \alpha_{n+1} \circ \phi_n(f)|| < \varepsilon_n, \tag{e 4.19}$$

for $f \in \mathcal{F}_n \cup \cup_{j=1}^{n-1} \phi_j(\mathcal{F}_j)$.

Then there is a *-homomorphism $\alpha\colon A \to A$ such that

$$\alpha(\phi_{l,\infty}(x)) = \lim_{i\to\infty} \phi_{i,\infty} \circ \mathrm{ad}(V_i) \circ \alpha_i \circ \phi_{l,i}(x), \tag{e 4.20}$$

$x \in A_l$ and $l = 1, 2, \ldots$, where $\phi_{l,i} = \phi_{i-1} \circ \phi_{i-2} \circ \cdots \circ \phi_l$. Moreover α is an automorphism.

It is also easy to check that

$$\alpha_{*i} = I(\psi^{\natural})_{*i}, \; i = 0, 1. \tag{e 4.21}$$

We also note that, by (e 4.19),

$$\alpha(\phi_{m,\infty}(f)) \approx_{1/2^{m-1}} \phi_{m,\infty} \circ \mathrm{ad}\, V_m \circ \alpha_m(f) \;\text{ for }\; f \in \mathcal{F}_m. \tag{e 4.22}$$

Denote by $j_n : C(X) \to C(X) \otimes M_{k(n)}$ the embdedding defined by $j_n(f) = f \otimes 1$.

Claim: for each $m \geq 1$ and $\tau \circ \phi_{m,\infty} \circ j_m(f) = \tau \circ \phi_{m,\infty} \circ j_m(f \circ \psi)$ for all $f \in C(X)$.

Fix $m \geq 1$. If $n \geq m + 1$, there is a projection $q_n \in M_{k(n)}$ such that

$$\phi_{m,n} \circ j_m(f) q_n = q_n \phi_{m,n} \circ j_m(f), \quad tr(q_n) = \frac{a_n}{b_n}$$

and, one may write that

$$\phi_{m,n} \circ j_m(f)(1 - q_n) = \mathrm{diag}(f(x(1, n)), f(x(2, n), ..., f(x(l(n), n)))$$

for all $f \in C(X)$. Therefore

$$\phi_{m,n} \circ j_m(f \circ \psi)(1 - q_n) = \mathrm{diag}(f(\psi(x(1, n))), f(\psi(x(2, n))), ..., f(\psi(x(l(n), n))))$$

for all $f \in C(X)$. Denote by $\bar{q}_n = \phi_{n,\infty}(q_n)$. For any $\varepsilon > 0$, and any $f \in C(X)$, by the virtue of Lemma 4.6, if n is sufficiently large, there is a permutation matrix $W_n \in M_{k(n)}$ such that

$$\|W_n^*[\phi_{m,n} \circ j_m(f)(1 - q_n)]W_n - \phi_{m,n} \circ j_m(f \circ \psi)(1 - q_n)\| < \varepsilon.$$

Put $\bar{W}_n = \phi_{n,\infty}(W_n)$. The above implies that

$$\|\bar{W}_n^* \phi_{m,\infty} \circ j_m(f)(1 - \bar{q}_n)\bar{W}_n - \phi_{m,\infty} \circ j_m(f \circ \psi)(1 - \bar{q}_n)\| < \varepsilon.$$

Thus we conclude that

$$|\tau(\phi_{m,\infty} \circ j_m(f)) - \tau(\phi_{m,\infty} \circ j_m(f \circ \psi))| < \frac{a_n}{b_n} + \varepsilon$$

for all sufficiently large n. Since $\lim_{n\to\infty} \frac{a_n}{b_n} = 0$. This implies that

$$\tau(\phi_{m,\infty} \circ j_m(f)) = \tau(\phi_{m,\infty} \circ j_m(f \circ \psi))$$

for all $f \in C(X)$. This proves the claim.

So

$$\tau \circ \phi_{m,\infty} \circ j_m(f) = \int_X f d\mu \tag{e 4.23}$$

for all $f \in C(X)$ and for some ψ-invariant Borel probability measure μ.

Now we verify that α has tracial Rokhlin property. For this end, we fix an integer $N \geq 1$, $\varepsilon > 0$ and a finite subset $\mathcal{F} \subset A$. For convenience, without loss of generality, we may assume that there exists $\mathcal{F}' \subset C(X) \otimes M_{k(m)}$ such that

$$\phi_{m,\infty}(\mathcal{F}') \supset \mathcal{F} \cup \cup_{j=1}^{N} \alpha^{-j}(\mathcal{F})$$

We may further assume that $\mathcal{F}'$ is in the unit ball of $C(X) \otimes M_{k(m)}$ for some integer $m \geq 1$. We may also assume that

$$\mathcal{F}' \subset \{(a_{i,j}) : a_{i,j} \in \mathcal{G}, 1 \leq i, j \leq k_{i_l}\},$$

where $\mathcal{G} \subset C(X)$ is a finite subset in the unit ball. Without loss of generality, we may further assume that

$$\mathcal{G} \subset \{f_1, f_2, ..., f_m\}.$$

Choose $\eta > 0$ such that, if $x, x' \in X$ and $\mathrm{dist}(x, x') < \eta$,

$$|f(x) - f(x')| < \frac{\varepsilon}{4N^2 8(k(m))^2} \tag{e 4.24}$$

for all $f \in \mathcal{G}$.

It follows from Lemma 4.7 that there are mutually disjoint open subsets $G_1, G_2, ..., G_L \subset X$ such that
(i) $\mathrm{diam}(G_i) < \eta$,
(ii) $\psi^j(\cup_{i=1}^{L} G) \cap \psi^l(\cup_{i=1}^{L} G_i)) = \emptyset$, if $j \neq l$ and $j, l = 0, 1, ..., N-1$,
(iii) $\mu(\partial(G_i)) = 0$ for all $\mu \in T_\psi$ and
(iv) $\mu(\cup_{j=0}^{N-1} \psi^j(\cup_{i=1}^{L} G_i)) > 1 - \varepsilon/16$ for all $\mu \in T_\psi$.

There is, for each i, another open subset S_i such that the closure of S_i is in G_i, and

$$\mu(S_i) > \mu(G_i) - \varepsilon/4LN \quad \text{and} \quad \mu(\partial(S_i)) = 0 \tag{e 4.25}$$

for all $\mu \in T_\psi$.

Let $g_i' \in C(X)$ such that $0 \leq g_i'(x) \leq 1$, $g_i'(x) = 0$ if $x \notin G_i$, $0 < g_i'(x)$ if $x \in G_i$ and $g_i'(x) = 1$ if $x \in \bar{S}_i$, $i = 1, 2, ..., L$. Define $a_i' \in C(X)$ such that $0 \leq a_i'(x) \leq 1$, $a_i'(x) > 0$ for all $x \in S_i$ and $a_i'(x) = 0$ if $x \notin S_i$, $i = 1, 2, ..., L$.

Fix $\eta_0 > 0$. Choose $m_1 \geq m$ such that $1/2^{m_1} < \min\{\varepsilon, \eta_0\}/16N^4$. We also assume that

$$\mathrm{dist}(a_i', \{f_1, f_2, ..., f_{m_1}\}) < \varepsilon/4N \quad \text{and} \quad \mathrm{dist}(g_i', \{f_1, f_2, ..., f_{m_1}\}) < \varepsilon/4N \tag{e 4.26}$$

$i = 1, 2, ..., L$.

We view g_i' and a_i' as elements in $E_{11} M_{k(m_1)} E_{11}$, where $\{E_{i,j}\}$ is a system of matrix unit for $M_{k(m_1)}$. Denote by $a_i'' = \phi_{m_1,m_1+1}(a_i')$ $a_i = \phi_{m_1,\infty}(a_i')$, $g_i'' = \phi_{m_1,m_1+1}(g_i')$ and $\bar{g}_i = \phi_{m_1,\infty}(g_i')$, $i = 1, 2, ..., L$.

Consider the hereditary C^*-subalgebra $\overline{a_i A a_i}$. Then, by the claim and by (e 4.23), there is a measure $\mu \in T_\psi$ such that

$$\mu(S_i) = k(m_1) \sup\{\tau(b) : 0 \leq b \leq 1, b \in \overline{a_i A a_i}\}, \ \ i = 1, 2, ..., L. \tag{e 4.27}$$

Since A has real rank zero, one obtains a projection $p_i \in \overline{a_iAa_i}$ such that

$$\tau(p_i) > \frac{\mu(S_i)}{k(m_1)} - \varepsilon/2LNk(m_1), \;\; i = 1, 2, ..., L \tag{e 4.28}$$

Put $\bar{E}_{i,j} = \phi_{m_1,\infty}(E_{i,j})$. Note $p_i \leq \bar{E}_{1,1}$. With $1_A = \sum_{i=1}^{k(m_1)} \bar{E}_{i,i}$, write

$$e_1 = \mathrm{diag}(\overbrace{\sum_{i=1}^{L} p_i, \sum_{i=1}^{L} p_i, \ldots, \sum_{i=1}^{L} p_i}^{k(m_1)}).$$

By the choice of η (see also (e 4.24) and Lemma 4.8—see also 3.2 of [17]), it follows that

$$\|e_1\phi_{m,\infty}(f) - \phi_{m,\infty}(f)e_1\| < \varepsilon/4N^2 \tag{e 4.29}$$

for all $f \in \mathcal{F}'$. By the definition of $\mathcal{F}'$, we have

$$\|\alpha^j(e_1)f - f\alpha^j(e_1)\| < \varepsilon/4N^2 \tag{e 4.30}$$

for $0 \leq j \leq N$ and all $f \in \mathcal{F}$.

Put $h_j = \phi_{m_1,\infty}(\mathrm{diag}(\overbrace{\sum_{i=1}^{L} g_i' \circ \psi^{j-1}, \sum_{i=1}^{L} g_i' \circ \psi^{j-1}, ..., \sum_{i=1}^{L} g_i' \circ \psi^{j-1}}^{k(m_1)}))$, $j = 1, 2, ..., N$. Note that

$$h_jh_i = h_ih_j = 0 \;\text{ if }\; i \neq j,\; i, j = 1, 2, ..., N. \tag{e 4.31}$$

We compute that

$$\mathrm{ad}\, V_{m_1}\alpha_{m_1}(h_1) = h_2. \tag{e 4.32}$$

Note that $e_1 \leq h_1$. It follows from (e 4.32) and (e 4.22) that

$$\|\alpha^j(h_1) - h_{j+1}\| < N(1/2^{m_1-1}) < 2N(1/2^{m_1}) < \min\{\varepsilon, \eta_0\}/8N^3, \tag{e 4.33}$$

$j = 1, 2, ..., N - 1$. Therefore, by (e 4.31),

$$\|\alpha^j(e_1)\alpha^i(e_1)\| < \min\{\varepsilon, \eta_0\}/4N^3, \;\; i \neq j,\; i, j = 0, 1, ..., N. \tag{e 4.34}$$

By choosing a small η_0 (which depends only on ε and N), it follows from [19, Lemma 2.5.6] that there are mutually orthogonal projections $e_1, e_2, ..., e_N$ such that

$$\|\alpha^j(e_1) - e_{j+1}\| < \varepsilon/N, \;\; j = 0, 1, ..., N - 1. \tag{e 4.35}$$

It follows that

$$\|\alpha(e_i) - e_{i+1}\| < \varepsilon, \;\; i = 1, 2, ..., N - 1 \tag{e 4.36}$$

By (e 4.30) and (e 4.35), we also have

$$\|e_if - fe_i\| < \varepsilon, \;\; i = 1, 2, ..., N \tag{e 4.37}$$

for all $f \in \mathcal{F}$.

By (e 4.28),

$$\begin{aligned}\tau(e_1) &= k(m_1)\tau(\sum_{i=1}^{L} p_i) > \sum_{i=1}^{L}(\mu(S_i) - \frac{\varepsilon}{2LN}) \\ &= \sum_{i=1}^{L}\mu(S_i) - \frac{\varepsilon}{2N} > \sum_{i=1}^{L}(\mu(G_i) - \frac{\varepsilon}{4LN}) - \frac{\varepsilon}{2N} \\ &= \mu(\cup_{i=1}^{L} G_i) - \frac{3\varepsilon}{4N}.\end{aligned}$$

By (e 4.21),

$$\tau(\alpha^j(e_1)) = \tau(e_1), \;\; j = 1, 2, ..., N-1. \tag{e 4.38}$$

Finally, by (iv) and (e 4.25), we compute that

$$\begin{aligned}\tau(\sum_{i=1}^{N} e_i) &= N\tau(e_1) > N\mu(\cup_{i=1}^{L} G_i) - \frac{3\varepsilon}{4} \\ &= \mu(\cup_{j=0}^{N-1}\psi^j(\cup_{i=1}^{L} G_i)) - \frac{3\varepsilon}{4} \\ &> 1 - \frac{\varepsilon}{16} - \frac{3\varepsilon}{4} > 1 - \varepsilon.\end{aligned}$$

Hence α has the tracial Rokhlin property. By (e 4.21), $\alpha_{*0}|_D = \mathrm{id}_D$. Since $\rho_A(D) = \rho_A(K_0(A))$, it follows from [22, Theorem 3.16] that α has the tracial cyclic Rokhlin property. □

Remark 4.11. In the proof of Theorem 4.10, one could work on more general inductive limits. For example, instead of considering a single summand $M_{k(n)}(C(X))$, one can consider unital simple C^*-algebras which are inductive limits of C^*-algebras with form $\oplus_{i=1}^{l(n)} M_{r(i,n)}(C(X))$ with each partial maps having the same form as (e 4.17). By doing that, more general dimensional groups D and more complicated $K_0(A)$ can be obtained. However, we choose this much simple construction to shed some light on the relation of the non-commutative dynamical systems and commutative dynamical systems which could be buried by some more complicated combinatory argument.

It should also be pointed out that the resulted crossed product C^*-algebra $A \rtimes_\alpha \mathbb{Z}$ obtained from Theorem 4.10 has tracial rank zero, by Theorem 2.4, and $A \rtimes_\alpha \mathbb{Z}$ is a unital simple AH-algebra.

Remark 4.12. In the case that $X = S^1$, let $\theta \in \mathbb{R}\backslash\mathbb{Q}$. Consider the rotation $\phi\colon X \to X$ by $\phi(z) = (\exp 2\pi i\theta)z$. Then ϕ is minimal and uniquely ergodic. Theorem 4.10 gives automorphisms α on unital simple $A\mathbb{T}$-algebras with tracial cyclic Rokhlin property. A modification of the proof of Theorem 4.10 will give an automorphism with the tracial Rokhlin property on every unital simple $A\mathbb{T}$-algebra with real rank zero. But all of them are approximately inner. However, one can use these approximately inner automorphisms with Rokhlin property in the proof of Theorem 3.5 instead of applying a deeper result of N. C. Phillips ([31]). It should be noted that, in this case the crossed product algebras $A \rtimes_\alpha \mathbb{Z}$ are again a unital simple $A\mathbb{T}$ algebra of real rank zero by [25, Corollary 3.6] (and [24]).

Not all connected finite CW complex X have minimal homeomorphisms. One of the interesting examples is Furstenberg transformation on $\mathbb{T}^k$.

Example 4.13. Let $\theta \in \mathbb{R}\backslash\mathbb{Q}$, $d \in \mathbb{Z}\backslash\{0\}$, and let $g_j \colon \mathbb{T}^j \to \mathbb{T}$ be a continuous map, $j = 1, 2, ..., k-1$. The Furstenberg transformation is defined by

$$\phi(z_1, z_2, ..., z_k) = (z_1 e^{2\pi i\theta}, g_1(z_1)z_2, ..., g_{k-1}((z_1, z_2, ..., z_{k-1}))z_k)$$

$|z_j| = 1$, $j = 1, 2, ..., k$. Furstenberg showed that ϕ is minimal, if all g_j are non-trivial, namely, $g_j(z_1, \ldots, z_j) = z_j^{d_j} e^{2\pi i f_j(z_1)}$ for some continuous function $f_j : \mathbb{T} \to \mathbb{R}$ with $d_j \neq 0$, $j = 1, 2, ..., k-1$. Thus $\phi^{\natural}_{*1} \neq \mathrm{id}_{K_1(C(\mathbb{T}^k))}$. Therefore the automorphism α constructed from ψ on $A_{\mathbb{T}^k}$ is not approximately inner since $\alpha_{*1} \neq \mathrm{id}_{K_1(A_{\mathbb{T}^k})}$. (See for example [13].)

References

[1] L. G. Brown and G. P. Pedersen, *C*-algebras of real rank zero*, J. Funct. Anal. 99(1991), 131 - 149.

[2] A. Connes, *Outer conjugacy classes of automorphisms of factors*, Ann. Sci. Éc. Norm. Sup. Ser. 4, **8**(1975), 383–420.

[3] G. A. Elliott, On the classification of C*-algebras of real rank zero, J. reine angew Math. **4**43(1993), 179 - 219.

[4] G. A. Elliott and G. Gong, *On the classification of C^*-algebras of real rank zero, II*, Ann. of Math. **1**44(1996), 497 - 610.

[5] G. A. Elliott and G. Gong, *On inductive limit of matrix algebras over the two-torus*, Amer. J. Math. **1**18(1996), 263 - 290.

[6] K. R. Goodearl, *Notes on a class of simple C*-algebras with real rank zero*, Publications Mathematiques **3**6(1992), 637 - 654.

[7] G. Gong and H. Lin, *Almost multiplicative morphisms and K-theory*, Internat. J. Math. **11** (2000), 983–1000.

[8] R. H. Herman and A. Ocneanu, *Stability for integer actions on UHF C*-algebras*, J. Funct. Anal. **59**(1984), 132–144.

[9] P. R. Halmos and H. E. Vaughan, *The marriage problem*, Amer. J. Math. **72**(1950), 214–215.

[10] A. Iwanik, M. Lemańczyk, and D. Rudolph, *Absolutely continuous cocycles over irrational rotations*, Israel J. Math. **83**(1993), 73–95.

[11] M. Izumi, *The Rohlin property for automorphisms of C*-algebras*, pages 191–206 in: *Mathematical Physics in Mathematics and Physics (Siena, 2000)*, Fields Inst. Commun. vol. 30, Amer. Math. Soc., Providence RI, 2001.

[12] M. Izumi, *Finite group actions on C^*-algebras with the Rohlin property. I*, Duke Math. J. **122**(2004), no. 2, 233–280.

[13] R. Ji, *On the crossed product C*-algebras associated with Furstenberg transformations on Tori*, Dissertation, State University of New York at Stony Brook, 1986.

[14] A. Kishimoto, *The Rohlin property for automorphisms of UHF algebras*, J. reine angew. Math. **465**(1995), 183–196.

[15] A. Kishimoto, *Automorphisms of $A\mathbb{T}$-algebras with the Rohlin property*, J. Operator Theory **4**0(1988), 277 - 294.

[16] A. Kishimoto, *Unbounded derivations in AT algebras*, J. Funct. Anal. **160**(1998), 270–311.

[17] H. Lin. *Homomorphisms from C^*-algebras of continuous trace*, Math. Scand. **86** (2000), 249–272.

[18] H. Lin, *Classification of simple tracially AF C*-algebras*, Canad. J. Math. **5**3(2001), 161 - 194.

[19] H. Lin *An introduction to the classification of amenable C*-algebras*, 2001, World Scientific.

[20] H. Lin, *Simple C^*-algebras with unique tracial states and quantized topological spaces*, Acta Math. Sin. (Engl. Ser.) **18** (2002), 181–198.

[21] H. Lin, *Classification of simple C^*-algebras with tracial topological rank zero*, Duke Math. J. **1**25(2004), 91 - 119.

[22] H. Lin, *Classification of homomorphisms and dynamical systems*, Trans. Amer. Math. Soc., to appear, arXiv:math.OA/0401240(2004).

[23] H. Lin *Minimal homeomorphisms and approximate conjugacy in measure*, arXiv:math.OA/0501262.

[24] H. Lin, *The Rokhlin property for automorphisms on simple C^*-algebras*, for Proceedings of GPOTS 2005, preprint.

[25] H. Lin and H. Osaka, The Rokholin property and the tracial topological rank, J. Functional Anal. **2**18(2005), 475 - 494.

[26] E. Lindenstrauss and B. Weiss, *Mean topological dimension*, Israel Journal of Mathematics **1**15(2000), 1 - 24.

[27] H. Nakamura, *Aperiodic automorphisms of certain simple C^*-algebras*, Operator algebras and applications, 145–157, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.

[28] H. Osaka and N. C. Phillips, Stable and real rank for crossed products by automorphisms with the tracial Rokhlin property, to appear in Ergod. Th. Dynam. Sys. (arXiv:math.OA/0409168(2004)).

[29] H. Osaka and N. C. Phillips, Furstenberg transformations on irrational rotation algebras, to appear in Ergod. Th. Dynam. Sys. (arXiv:math.OA/0409169(2004)).

[30] N. C. Phillips, *Crossed products by finite cyclic group actions with the tracial Rokhlin property*, arXiv.org math.OA/0306410.

[31] N. C. Phillips, The tracial Rokhlin property is generic, in preparation.

[32] M. Rørdam, *Classification of certain infinite simple C^*-algebras*, J. Funct. Anal. **131** (1995), 415–458.

[33] K. Thomsen, On isomorphisms of inductive limit C*-algebras, Proc. A. M. S. **1**13(1991), 947 - 953.

[34] N. E. Wegge-Olsen, *K-theory and C^*-algebras. A friendly approach*, Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1993.